

MATH 31A DISCUSSION

JED YANG

1. MORE APPLICATIONS OF THE DERIVATIVE

1.1. Recall.

1.1.1. *Concavity.* If $f'(x)$ is increasing (or $f''(x) > 0$), then f is *concave up* at x . If $f'(x)$ is decreasing (or $f''(x) < 0$), then f is *concave down* at x .

1.1.2. *Inflection.* If $f''(c) = 0$ and $f''(x)$ changes sign at $x = c$, then $f(x)$ has a point of inflection at $x = c$.

1.1.3. *Second Derivative Test.* Let f be differentiable and c a critical point ($f'(c) = 0$).

- (a) If $f''(c) > 0$ then $f(c)$ is a local minimum.
- (b) If $f''(c) < 0$ then $f(c)$ is a local maximum.
- (c) If $f''(c) = 0$ then it is inconclusive, $f(c)$ may be a local min, max, or neither.

1.2. **Exercise 4.4.24.** Find the critical points of $f(x) = \sin^2 x + \cos x$, x in $[0, \pi]$, and use the Second Derivative Test to determine whether each corresponds to a local minimum or maximum.

Solution. Notice $f'(x) = 2 \sin x \cos x - \sin x$. Setting $f'(x) = 0$ and solving, we get $\sin x(2 \cos x - 1) = 0$, so $x = 0, \pi/3, \pi$. Now $f''(x) = 2 \cos^2 x - 2 \sin^2 x - \cos x$. So $f''(0) = 1$, $f''(\pi/3) = -\frac{3}{2}$, $f''(\pi) = 3$, yielding local minima at $x = 0, \pi$ and maximum at $x = \pi/3$. \square

1.3. **Exercise 4.4.53.** If $f'(c) = 0$ and $f(c)$ is neither a local min or max, must $x = c$ be a point of inflection? This is true of most “reasonable” examples, but it is not true in general. Let

$$f(x) = \begin{cases} x^2 \sin \frac{1}{x} & \text{for } x \neq 0, \\ 0 & \text{for } x = 0. \end{cases}$$

- Use the limit definition of the derivative to show that $f'(0)$ exists and $f'(0) = 0$.
- Show that $f(0)$ is neither a local min nor max.
- Show that $f'(x)$ changes sign infinitely often near $x = 0$ and conclude that $f(x)$ does not have a point of inflection at $x = 0$.

Solution. Recall Exercise 3.7.92 from 10/20.

Recall $f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$. So by definition, $f'(0) = \lim_{h \rightarrow 0} \frac{f(h) - f(0)}{h} = \lim_{h \rightarrow 0} h \sin \frac{1}{h}$. Using Squeeze Theorem and $-|h| \leq h \sin \frac{1}{h} \leq |h|$, we get that $f'(0) = 0$. Away from $x = 0$, we can use the formula and get $f'(x) = 2x \sin \frac{1}{x} + x^2 \cos \frac{1}{x} \cdot (-1) \frac{1}{x^2} = 2x \sin \frac{1}{x} - \cos \frac{1}{x}$. Now $\lim_{x \rightarrow 0} f'(x)$ does not exist since $\lim_{x \rightarrow 0} 2x \sin \frac{1}{x} = 0$ by Squeeze Theorem but $\lim_{x \rightarrow 0} \cos \frac{1}{x}$ does not exist. \square

1.4. **Exercise 4.5.59.**

1.5. **Exercise 4.5.68.**

1.6. **Bonus Question.** Assume $f''(x)$ exists and $f''(x) > 0$ for all x . Show that $f(x)$ cannot be always negative.

Solution. If $f'(x) \equiv 0$, then $f''(x) \equiv 0$, a contradiction. So there exists b such that $f'(b) \neq 0$. Consider the tangent line at $x = b$ to $f(x)$. It is given by the equation $y = f'(b)(x - b) + f(b)$. Consider $g(x) = f(x) - f'(b)(x - b) - f(b)$. Notice that $g'(x) = f'(x) - f'(b)$ and $g''(x) = f''(x)$. So $g(b) = g'(b) = 0$, and $g''(x) > 0$ for all x . Hence $g'(x)$ is increasing. In particular, $g'(x) < 0$ for $x < b$ and $g'(x) > 0$ for $x > b$. If $x > b$, then by MVT, we get

$$\frac{g(x) - g(b)}{x - b} = g'(c)$$

for some c in the interval (b, x) . In otherwords, since $c > b$, we have that $g'(c) > 0$ and $x - b > 0$, hence $g(x) - g(b) > 0$. Similarly, if $x < b$, we get $g'(c) < 0$, $x - b < 0$, so $g(x) - g(b) > 0$ as well. We thus conclude $g(x) \geq g(b)$ for all x . So $f(x) \geq f'(b)(x - b) + f(b)$ for all x . Since $f'(b) \neq 0$, there exists x far enough from the origin such that $f'(b)(x - b) + f(b) > 0$, as desired. \square