

Math 4990 Scissors congruence and Dehn invariants

JED YANG

§1. In \mathbb{R}^2 .

Theorem 1. *If P and Q are polygons with the same area, they are scissors congruent.*

Proof. Triangulate. Triangles are scissors congruent to rectangles. Rectangles (of the same area) are scissors congruent to each other. \square

§2. \mathbb{Q} -vector spaces. For $M = \{m_1, \dots, m_k\} \subseteq \mathbb{R}$, let

$$V(M) := \left\{ \sum_{i=1}^k q_i m_i : q_i \in \mathbb{Q} \right\} \subseteq \mathbb{R}$$

denote the set of all linear combinations of numbers in M with rational coefficients.

Note that $V(M)$ is a finite-dimensional vector space over the field \mathbb{Q} .

The dimension is the size of any minimal generating set. As M generates,

$$\dim_{\mathbb{Q}} V(M) \leq k = |M|.$$

Considering \mathbb{Q} as a 1-dimensional vector space over itself, a linear map

$$f : V(M) \rightarrow \mathbb{Q}$$

of \mathbb{Q} -vector spaces is a \mathbb{Q} -linear function, which satisfies these properties:

- (i) $f(x) + f(y) = f(x + y)$ for $x, y \in V(M)$
- (ii) $f(qx) = qf(x)$ for $x \in V(M)$, $q \in \mathbb{Q}$

§3. Dehn invariants. Let P be a 3-dimensional polyhedron. For an edge e of P , let $\ell(e)$ denote its length, and $\phi(e)$ its **dihedral angle**, defined as the angle between the two faces meeting at e .

Let M_P be the set of all dihedral angles of P and π .

Cube C has $M_C = \{\pi/2, \pi\}$.

For a \mathbb{Q} -linear function $f : V(M) \rightarrow \mathbb{Q}$ with $f(\pi) = 0$, define the **Dehn invariant** $D_f(P)$ of P (with respect to f) by

$$D_f(P) := \sum_{e \in P} \ell(e) \cdot f(\phi(e)),$$

where the sum runs over all edges e of the polyhedron P .

For any such f , as $f(\pi/2) = \frac{1}{2}f(\pi) = 0$, we have $D_f(C) = 0$.

§4. In \mathbb{R}^3 .

Theorem 2 (Sydler 1.67). *If P and Q are not scissors congruent, then there is some f such that the Dehn invariant $D_f(P) \neq D_f(Q)$.*

We will not prove Sydler theorem, but we will prove Dehn–Hadwiger theorem next time, which establishes the converse.

§5. Irrationality.

Theorem 3. For odd integer $n \geq 3$,

$$\frac{1}{\pi} \arccos \frac{1}{\sqrt{n}}$$

is irrational.

Proof. Recall addition formula

$$\cos(\alpha \pm \beta) = \cos \alpha \cos \beta \mp \sin \alpha \sin \beta.$$

Summing yields

$$\cos(\alpha + \beta) + \cos(\alpha - \beta) = 2 \cos \alpha \cos \beta.$$

Let $\varphi_n = \arccos \frac{1}{\sqrt{n}}$, so $\cos \varphi_n = \frac{1}{\sqrt{n}}$. Substituting $\alpha = k\varphi_n$ and $\beta = \varphi_n$ yields

$$\cos(k+1)\varphi_n + \cos(k-1)\varphi_n = 2 \cos \varphi_n \cos k\varphi_n$$

Claim 3.1. For all integers $k \geq 0$,

$$\cos k\varphi_n = \frac{A_k}{\sqrt{n}^k}$$

for some integer $A_k \in \mathbb{Z}$ such that $n \nmid A_k$.

Proof. Indeed, $A_0 = A_1 = 1$. By induction, we have

$$\begin{aligned} \cos(k+1)\varphi_n &= 2 \cos \varphi_n \cos k\varphi_n - \cos(k-1)\varphi_n \\ &= 2 \frac{1}{\sqrt{n}} \frac{A_k}{\sqrt{n}^k} - \frac{A_{k-1}}{\sqrt{n}^{k-1}} = \frac{2A_k - nA_{k-1}}{\sqrt{n}^{k+1}}, \end{aligned}$$

so

$$A_{k+1} = 2A_k - nA_{k-1}$$

is an integer. Moreover, if $n \mid A_{k+1}$ then $n \mid 2A_k$. But $n \geq 3$ is odd and $n \nmid A_k$, a contradiction. \square

Suppose, towards a contradiction, that

$$\frac{1}{\pi} \arccos \frac{1}{\sqrt{n}} = \frac{p}{q}$$

with integers $p, q > 0$, then taking cosine of both sides of

$$q\varphi_n = p\pi$$

yields

$$\pm 1 = \cos p\pi = \cos q\varphi_n = \frac{A_q}{\sqrt{n}^q},$$

and hence $\sqrt{n}^q = \pm A_q$ is an integer. In fact, $q > 1$ as $0 < \arccos \frac{1}{\sqrt{n}} < \pi/2$. So $q \geq 2$ and $n \mid \sqrt{n}^q \mid A_q$, a contradiction. \square

2014.09.23

§6. Proof of Dehn–Hadwiger theorem.

Theorem 4 (Dehn–Hadwiger 1.62). *Let P be a polyhedron decomposed into finitely many polyhedral pieces P_1, \dots, P_k . Let $f : V(M) \rightarrow \mathbb{Q}$ be a d -function, where $M \subset \mathbb{R}^3$ is finite and*

$$M \supseteq M_P \cup M_{P_1} \cup \dots \cup M_{P_k}.$$

Then

$$D_f(P) = \sum_{i=1}^k D_f(P_i).$$

Proof. Let S denote all edge segments. For a polyhedron Q and a straight line segment s , define the dihedral angle as follows. If s is part of an edge e , then it shares the dihedral angle of e . If s lies on a face or in the interior, then the dihedral angle is π or 2π , respectively. Otherwise, say it is 0.

Let $\phi(s)$ and $\phi_i(s)$ denote the dihedral angle of s with respect to P and P_i , respectively. The key observation is that

$$\phi(s) = \sum_i \phi_i(s), \tag{*}$$

for any $s \in S$, regardless of its spatial relationships to P and the P_i .

Dehn invariants can be calculated over (all) edge segments:

$$\begin{aligned} D_f(Q) &= \sum_{e \in E(Q)} \ell(e) \cdot f(\phi(e)) \\ &= \sum_{e \in E(Q)} f(\phi(e)) \cdot \sum_{\substack{s \in S \\ s \subseteq e}} \ell(s) \\ &= \sum_{e \in E(Q)} \sum_{\substack{s \in S \\ s \subseteq e}} \ell(s) f(\phi(s)) \\ &= \sum_{s \in S} \ell(s) f(\phi(s)), \end{aligned} \tag{†}$$

where ϕ is with respect to Q .

The rest is a simple calculation:

$$\begin{aligned}
\sum_i D_f(P_i) &= \sum_i \sum_{e \in E(P_i)} \ell(e) \cdot f(\phi_i(e)) \\
&= \sum_i \sum_{s \in S} \ell(s) \cdot f(\phi_i(s)) && \text{by } (\dagger) \\
&= \sum_{s \in S} \sum_i \ell(s) \cdot f(\phi_i(s)) \\
&= \sum_{s \in S} \ell(s) \cdot \sum_i f(\phi_i(s)) \\
&= \sum_{s \in S} \ell(s) \cdot f\left(\sum_i \phi_i(s)\right) && \text{by } \mathbb{Q}\text{-linearity} \\
&= \sum_{s \in S} \ell(s) \cdot f(\phi(s)) && \text{by } (*) \\
&= D_f(P), && \text{by } (\dagger)
\end{aligned}$$

as desired. □

Corollary 5 (1.62). *Let P and Q be two polyhedra, and $M \subset \mathbb{R}$ finite such that $M \supseteq M_P \cup M_Q$. If $f : V(M) \rightarrow \mathbb{Q}$ is any \mathbb{Q} -linear function with $f(\pi) = 0$ such that $D_f(P) \neq D_f(Q)$, then P and Q are not scissors congruent.*

Proof. Suppose P and Q are scissors congruent, and let $(M$ and) f be given.

Fix some common decomposition: P is decomposed into P_1, \dots, P_k and Q is decomposed into Q_1, \dots, Q_k , where P_i and Q_i are congruent. Let $M' \supseteq M$ be a finite set that includes all dihedral angles that appear.

Extend f to $f' : V(M') \rightarrow \mathbb{Q}$ by specifying the values of f' on new basis elements (and keep the old ones the same). Then

$$D_f(P) = D_{f'}(P) = \sum_{i=1}^k D_{f'}(P_i) = \sum_{i=1}^k D_{f'}(Q_i) = D_{f'}(Q) = D_f(Q),$$

as desired. □